

## No sex difference in mouse digit ratio: reply to Voracek

By: P. L. Hurd and [Douglas Wahlsten](#)

Hurd, P.L., & Wahlsten, D. (2006) No sex difference in mouse digit ratio: reply to Voracek. *Genes, Brain & Behavior*, 5: 300-302.

Made available courtesy of Wiley-Blackwell: The definitive version is available at

<http://www3.interscience.wiley.com>

**\*\*\*Reprinted with permission. No further reproduction is authorized without written permission from Wiley-Blackwell. This version of the document is not the version of record. Figures and/or pictures may be missing from this format of the document.\*\*\***

### **Article:**

Voracek specializes in the study of human non-neural morphology (Voracek & Fisher 2002) and suicide (Voracek 2005), and he does not appear to appreciate the methods used in experimental mouse genetics. We (Bailey et al. 2005) investigated 2D:4D digit ratio in inbred mouse strains, because the findings can lead directly and fruitfully to a genetic analysis of factors that lead to higher and lower ratios (we follow the convention of using 2D:4D to refer to this digit ratio as have all previous published animal and human studies of hind limb digit ratios. We know of no published work using Voracek's 2T:4T nomenclature. A PubMed search over the past 5 years uncovered not one paper using 2T:4T but 135 using 2D:4D, including all seven investigating hind limb digits). Indeed, our study detected substantial and interesting strain differences, although significant sex differences were not found. Voracek is evidently disappointed at the lack of a sex difference in our data and seeks to dismiss them by branding our work a 'pilot study' whose results 'diverge from other mouse evidence'.

The sum total of the 'mouse evidence' for the genus *Mus* is three studies. Brown et al. (2002) measured left and right hind paws of 32 weanling and 39 adult 'outbred lab mice' of unspecified origin, and they found a significantly lower 2D:4D ratio for males than females only for the right paw. Manning et al. (2003) measured only the left hind paw for 111 mice of unspecified age and strain. The poorly specified genetic composition of the two populations is appalling. Manning et al. did not even replicate the finding of Brown et al. for the right paw; hence, it is difficult to see how our data for a larger sample of 175 inbred mice from eight well-defined and easily replicable strains could 'diverge' from those two. Furthermore, for three strains with lower average digit ratios (BALB/cByJ, BTBR/T + tf/J and C3H/HeJ), we did find a lower ratio for males than females, although the sex difference was too small to achieve significance. While differences in measurement techniques (see discussion in Bailey et al. 2005) prevent rigorous comparisons of ratio values across studies, note that the absolute values from the Manning et al. study are almost exactly the same as the absolute values and sex differences of the C3H/HeJ strain in our study.

Voracek suggests that the evidence supporting a correlation between finger length ratio and developmental androgen exposure is very strong. We regard the evidence as circumstantial but suggestive. The Lutchmaya et al. (2004) study found a correlation between testosterone : estradiol and 2D:4D ratios in a small, combined sample of 18 males and 15 females. Differences in 2D:4D between human ethnic groups dwarf differences between the sexes, and sex differences also vary considerably across ethnic groups (Manning et al. 2000; McFadden et al. 2005). This suggests that the testosterone exposure story is at least an over-simplification. The only experimental study that manipulated developmental testosterone was Romano et al. (2005), which produced a change in hind limb digit ratio in pheasants.

More data are needed to illuminate the true relationship between developmental variables and digit ratio. For example, it would be interesting to study an F2 hybrid population, because extreme inbreeding might disrupt the regulation of developmental processes involving the digits. For the average effect size of the sex difference that

we observed ( $d = 0.04$ ), the research would need to be done with samples of at least 5000 per sex to achieve power of 90% when using Type I error probability = 0.05 (Wahlsten 1991).

## References

- Bailey, A.A., Wahlsten, D. & Hurd, P.L. (2005) Digit ratio (2D:4D) and behavioral differences between inbred mouse strains. *Genes Brain Behav* 4, 318–323.
- Brown, W.M., Finn, C.J. & Breedlove, S.M. (2002) Sexual dimorphism in digit-length ratios of laboratory mice. *Anat Rec* 267, 231–234.
- Burley, N.T. & Foster, V.S. (2004) Digit ratio varies with sex, egg order and strength of mate preference in zebra finches. *Proc R Soc Lond B Biol Sci* 271, 239–244.
- Cummins, H. & Midlo, C. (1961) *Finger Prints, Palms and Soles: An Introduction to Dermatoglyphics*. Dover, New York.
- Durham, N.M., Fox, K.M. & Plato, C.C., eds. (2000) *The State of Dermatoglyphics: The Science of Finger and Palm Prints*. Edwin Mellen Press, Lewiston, NY.
- Fink, B., Neave, N. & Manning, J.T. (2003) Second to fourth digit ratio, body mass index, waist-to-hip ratio, and waist-to-chest ratio: their relationships in heterosexual men and women. *Ann Hum Biol* 30, 728–738.
- Fitch, R.H. & Denenberg, V.H. (1998) A role for ovarian hormones in sexual differentiation of the brain. *Behav Brain Sci* 21, 311–352.
- Forstmeier, W. (2005) Quantitative genetics and behavioural correlates of digit ratio in the zebra finch. *Proc R Soc Lond B Biol Sci* 272, 2641–2649.
- Lutchmaya, S., Baron-Cohen, S., Raggatt, P., Knickmeyer, R. & Manning, J.T. (2004) 2nd to 4th digit ratios, fetal testosterone and estradiol. *Early Hum Dev* 77, 23–28.
- Manning, J.T. (2002) *Digit Ratio: a Pointer to Fertility, Behavior, and Health*. Rutgers University Press, New Brunswick, NJ.
- Manning, J.T., Barley, L., Walton, J., Lewis-Jones, D., Trivers, R.L., Singh, D., Thornhill, R., Rohde, P., Bereckei, T., Henzi, P., Soler, M. & Sved, A. (2000) The 2nd:4th digit ratio, sexual dimorphism, population differences, and reproductive success: evidence for sexually antagonistic genes. *Evol Hum Behav* 21, 163–183.
- Manning, J.T., Callow, M. & Bundred, P.E. (2003) Finger and toe ratios in humans and mice: implications for the aetiology of diseases influenced by HOX genes. *Med Hypotheses* 60, 340–343.
- McFadden, D. & Bracht, M.S. (2003) The relative lengths and weights of metacarpals and metatarsals in baboons (*Papio hamadryas*). *Horm Behav* 43, 347–355.
- McFadden, D. & Bracht, M.S. (2005) Sex differences in the relative lengths of metacarpals and metatarsals in gorillas and chimpanzees. *Horm Behav* 47, 99–111.
- McFadden, D. & Shubel, E. (2002) Relative lengths of fingers and toes in human males and females. *Horm Behav* 42, 492–500.
- McFadden, D., Loehlin, J.C., Breedlove, S.M., Lippa, R.A., Manning, J.T. & Rahman, Q. (2005) A reanalysis of five studies on sexual orientation and the relative length of the 2nd and 4th fingers (the 2D:4D ratio). *Arch Sex Behav* 34, 341–356.
- McMechan, A.P., O’Leary-Moore, S.K., Morrison, S.D. & Hannigan, J.H. (2004) Effects of prenatal alcohol exposure on forepaw digit length and digit ratios in rats. *Dev Psychobiol* 34, 251–258.
- Putz, D.A., Gaulin, S.J.C., Sporter, R.J. & McBurney, D.H. (2004) Sex hormones and finger length: what does 2D:4D indicate? *Evol Hum Behav* 25, 182–199.
- Romano, M., Rubolini, D., Martinelli, R., Alquati, A.B. & Saino, M. (2005) Experimental manipulation of yolk testosterone affects digit length ratios in the ring-necked pheasant (*Phasianus colchicus*). *Horm Behav* 48, 342–346.
- Roney, J.R., Whitham, J.C., Leoni, M., Bellem, A., Wielebnowski, N. & Maestripieri, D. (2004) Relative digit lengths and testosterone levels in Guinea baboons. *Horm Behav* 45, 285–290.
- Sorenson Jamison, C., Meier, R.J. & Campbell, B.C. (1993) Dermatoglyphic asymmetry and testosterone levels in normal males. *Am J Phys Anthropol* 90, 185–198.
- Voracek, M. (2005) The social ecology of intelligence and suicide in Belarus. *J Soc Psychol* 145, 613–617.
- Voracek, M. & Fisher, M.L. (2002) Shapely centrefolds? Temporal change in body measures: trend analysis. *BMJ* 325, 1447–1448.
- Wahlsten, D. (1991) Sample size to detect a planned contrast and a one degree-of-freedom interaction effect. *Psychol Bull* 110, 587–595.